

Some bialgebroids constructed by Kadison and Connes-Moscovici are isomorphic *

Florin Panaite

Institute of Mathematics of the Romanian Academy
PO-Box 1-764, RO-014700 Bucharest, Romania
e-mail: Florin.Panaite@imar.ro

Freddy Van Oystaeyen

Department of Mathematics and Computer Science
University of Antwerp, Middelheimlaan 1
B-2020 Antwerp, Belgium
e-mail: Francine.Schoeters@ua.ac.be

Abstract

We prove that a certain bialgebroid introduced recently by Kadison is isomorphic to a bialgebroid introduced earlier by Connes and Moscovici. At the level of total algebras, the isomorphism is a consequence of the general fact that an L-R-smash product over a Hopf algebra is isomorphic to a diagonal crossed product.

1 Introduction

Let H be a Hopf algebra with bijective antipode, A a left H -module algebra and denote as usual $A^e = A \otimes A^{op}$. In [10], Kadison constructed a bialgebroid (in the sense of [12]) with base A and having as total algebra a certain algebra structure on $A^e \otimes H$, which we denote by $A^e \diamond H$. We first note that this algebra structure is actually an L-R-smash product $A^e \bowtie H$ as introduced in [14]. An L-R-smash product is isomorphic to a diagonal crossed product as in [9], [6], hence $A^e \bowtie H \simeq A^e \bowtie H$. Finally, the diagonal crossed product $A^e \bowtie H$ is isomorphic to a certain algebra, which we denote by $A \odot H \odot A$, used by Connes and Moscovici in [8] (see also [11]) and which is also a bialgebroid over A . It turns out that the resulting algebra isomorphism is actually a bialgebroid isomorphism between $A^e \diamond H$ and $A \odot H \odot A$.

We establish also a universal property of $A^e \diamond H$ as a bialgebroid and we give an example (up to isomorphism) of the type $A \odot H \odot A$: the Cibils-Rosso algebra from [7].

2 The isomorphism

We work over a ground field k . All algebras, linear spaces etc. will be over k ; unadorned $\otimes$ means $\otimes_k$. Throughout H will be a Hopf algebra with bijective antipode S . We use the following version of Sweedler's Σ -notation: $\Delta(h) = h_1 \otimes h_2$ for $h \in H$.

*Research partially supported by the EC programme LIEGRITS, RTN 2003, 505078, and by the project "New techniques in Hopf algebras and graded ring theory" of the Flemish and Romanian Ministries of Research.

Recall that the L-R-smash product over a cocommutative Hopf algebra was introduced in [1], [2], [4], [5], and generalized to an arbitrary Hopf algebra in [14] as follows: if $\mathcal{A}$ is an H -bimodule algebra, the L-R-smash product $\mathcal{A} \bowtie H$ is the following algebra structure on $\mathcal{A} \otimes H$:

$$(\varphi \bowtie h)(\varphi' \bowtie h') = (\varphi \cdot h'_2)(h_1 \cdot \varphi') \bowtie h_2 h'_1, \quad \forall \varphi, \varphi' \in \mathcal{A}, h, h' \in H.$$

The diagonal crossed product $\mathcal{A} \bowtie H$ is the following algebra structure on $\mathcal{A} \otimes H$, see [9], [6]:

$$(\varphi \bowtie h)(\varphi' \bowtie h') = \varphi(h_1 \cdot \varphi' \cdot S^{-1}(h_3)) \bowtie h_2 h', \quad \forall \varphi, \varphi' \in \mathcal{A}, h, h' \in H.$$

Proposition 2.1 ([14]) $\mathcal{A} \bowtie H \simeq \mathcal{A} \bowtie H$ as algebras, the isomorphism being given as follows:

$$\begin{aligned} \nu : \mathcal{A} \bowtie H &\rightarrow \mathcal{A} \bowtie H, & \nu(\varphi \bowtie h) &= \varphi \cdot h_2 \bowtie h_1, \\ \nu^{-1} : \mathcal{A} \bowtie H &\rightarrow \mathcal{A} \bowtie H, & \nu^{-1}(\varphi \bowtie h) &= \varphi \cdot S^{-1}(h_2) \bowtie h_1. \end{aligned}$$

Let now A be a left H -module algebra, denote by A^{op} the opposite algebra and by $A^e = A \otimes A^{op}$. The bialgebroid introduced by Kadison in [10] has as total algebra the following algebra structure on $A^e \otimes H$, which we denote by $A^e \diamond H$:

$$(a \otimes b \otimes h)(a' \otimes b' \otimes h') = a(h_1 \cdot a') \otimes b'(S(h'_2) \cdot b) \otimes h_2 h'_1,$$

for all $a, a', b, b' \in A$ and $h, h' \in H$, where the multiplication on the second component is made in A (not in A^{op}). On the other hand, the opposite algebra A^{op} becomes a right H -module algebra with action $a \cdot h = S(h) \cdot a$, hence $A^e = A \otimes A^{op}$ is an H -bimodule algebra with actions $h \cdot (a \otimes b) \cdot h' = h \cdot a \otimes b \cdot h'$, for all $a, b \in A$ and $h, h' \in H$, so we may consider the L-R-smash product $A^e \bowtie H$.

Proposition 2.2 $A^e \diamond H = A^e \bowtie H$.

The bialgebroid introduced by Connes and Moscovici in [8] and further studied in [11] has as total algebra the following algebra structure on $A \otimes H \otimes A$, which we denote by $A \odot H \odot A$:

$$(a \otimes h \otimes b)(a' \otimes h' \otimes b') = a(h_1 \cdot a') \otimes h_2 h' \otimes (h_3 \cdot b')b,$$

for all $a, a', b, b' \in A$ and $h, h' \in H$. We may consider again the H -bimodule algebra A^e as above and the diagonal crossed product $A^e \bowtie H$.

Proposition 2.3 We have an algebra isomorphism

$$A \odot H \odot A \simeq (A \otimes A^{op}) \bowtie H, \quad a \otimes h \otimes b \mapsto (a \otimes b) \bowtie h.$$

Corollary 2.4 $A^e \diamond H \simeq A \odot H \odot A$ as algebras, an explicit isomorphism being given by

$$\begin{aligned} A^e \diamond H &\rightarrow A \odot H \odot A, & a \otimes b \otimes h &\mapsto a \otimes h_1 \otimes h_2 \cdot b, \\ A \odot H \odot A &\rightarrow A^e \diamond H, & a \otimes h \otimes b &\mapsto a \otimes S(h_2) \cdot b \otimes h_1. \end{aligned}$$

We recall from [10], [8], [11] the structure of $A^e \diamond H$ and $A \odot H \odot A$ as bialgebroids over A :

- (1) source maps: $A \rightarrow A^e \diamond H$, $a \mapsto (a \otimes 1) \otimes 1$ and $A \rightarrow A \odot H \odot A$, $a \mapsto a \otimes 1 \otimes 1$;
- (2) target maps: $A \rightarrow A^e \diamond H$, $a \mapsto (1 \otimes a) \otimes 1$ and $A \rightarrow A \odot H \odot A$, $a \mapsto 1 \otimes 1 \otimes a$;
- (3) comultiplications:

$$\begin{aligned} A^e \diamond H &\rightarrow (A^e \diamond H) \otimes_A (A^e \diamond H), \quad (a \otimes b) \otimes h \mapsto ((a \otimes 1) \otimes h_1) \otimes_A ((1 \otimes b) \otimes h_2), \\ A \odot H \odot A &\rightarrow (A \odot H \odot A) \otimes_A (A \odot H \odot A), \quad a \otimes h \otimes b \mapsto (a \otimes h_1 \otimes 1) \otimes_A (1 \otimes h_2 \otimes b); \end{aligned}$$

- (4) counits:

$$\begin{aligned} \varepsilon : A^e \diamond H &\rightarrow A, \quad \varepsilon((a \otimes b) \otimes h) = a(h \cdot b), \\ \varepsilon : A \odot H \odot A &\rightarrow A, \quad \varepsilon(a \otimes h \otimes b) = a\varepsilon(h)b. \end{aligned}$$

Theorem 2.5 *$A^e \diamond H$ and $A \odot H \odot A$ are isomorphic as bialgebroids over A , where the isomorphism is given as in Corollary 2.4 at the level of total algebras and as identity at the level of base algebras.*

Remark 2.6 Assume that H is involutive, i.e. $S^2 = id$. In this case $A^e \diamond H$ and $A \odot H \odot A$ have antipodes making them Hopf algebroids (in the sense of [3]), namely (see [10], [11]):

$$\begin{aligned} A^e \diamond H &\rightarrow A^e \diamond H, \quad (a \otimes b) \otimes h \mapsto (b \otimes a) \otimes S(h), \\ A \odot H \odot A &\rightarrow A \odot H \odot A, \quad a \otimes h \otimes b \mapsto S(h_3) \cdot b \otimes S(h_2) \otimes S(h_1) \cdot a. \end{aligned}$$

Then one can see that the isomorphism from Theorem 2.5 commutes with these antipodes, i.e. it is a *strict* isomorphism of Hopf algebroids (in the terminology of [3]).

Example 2.7 Assume that H is finite dimensional. Obviously H^* , the linear dual of H , is a left $H \otimes H^{op}$ -module algebra, with action $(h \otimes h') \cdot f = h \rightharpoonup f \leftharpoonup h'$, for all $h, h' \in H$ and $f \in H^*$, where $\rightharpoonup$ and $\leftharpoonup$ are the regular actions of H on H^* given by $(h \rightharpoonup f)(h') = f(h'h)$ and $(f \leftharpoonup h')(h) = f(h'h)$. Hence, we can consider the bialgebroid $H^* \odot (H \otimes H^{op}) \odot H^*$ over H^* ; by Proposition 2.3 we have $H^* \odot (H \otimes H^{op}) \odot H^* \simeq (H^* \otimes H^{*op}) \bowtie (H \otimes H^{op})$ as algebras. The algebra $Z = (H^* \otimes H^{*op}) \bowtie (H \otimes H^{op})$ was considered before in [13], it is isomorphic to the Cibils-Rosso algebra X from [7] having the property that modules over it coincide with H^* -Hopf bimodules. Consequently, we obtain that the Cibils-Rosso algebra is a bialgebroid over H^* .

3 A universal property of $A^e \diamond H$ as bialgebroid

Kadison found in [10] a universal property of $A^e \diamond H$ as algebra. We first establish an equivalent formulation, emphasizing the presence of the algebra $A \otimes A^{op}$:

Proposition 3.1 *Denote by $i : A \otimes A^{op} \rightarrow (A \otimes A^{op}) \diamond H$, $i(a \otimes b) = (a \otimes b) \otimes 1$ and $j : H \rightarrow (A \otimes A^{op}) \diamond H$, $j(h) = (1 \otimes 1) \otimes h$, the canonical algebra inclusions. If R is an associative algebra and $u : A \otimes A^{op} \rightarrow R$ and $v : H \rightarrow R$ are algebra maps such that*

$$v(h_1)u(a \otimes S(h_2) \cdot b) = u(h_1 \cdot a \otimes b)v(h_2),$$

for all $a, b \in A$ and $h \in H$, there exists a unique algebra map $\omega : (A \otimes A^{op}) \diamond H \rightarrow R$ such that $\omega \circ i = u$ and $\omega \circ j = v$. The map ω is given by $\omega((a \otimes b) \otimes h) = u(a \otimes h_2 \cdot b)v(h_1)$.

Recall from [12] that $A \otimes A^{op}$ is a bialgebroid over A , with source map $\alpha : A \rightarrow A \otimes A^{op}$, $a \mapsto a \otimes 1$, target map $\beta : A \rightarrow A \otimes A^{op}$, $a \mapsto 1 \otimes a$, coproduct $\Delta : A \otimes A^{op} \rightarrow (A \otimes A^{op}) \otimes_A (A \otimes A^{op})$, $\Delta(a \otimes b) = (a \otimes 1) \otimes_A (1 \otimes b)$ and counit $\varepsilon : A \otimes A^{op} \rightarrow A$, $\varepsilon(a \otimes b) = ab$. With respect to this bialgebroid structure on $A \otimes A^{op}$, we have a universal property of $(A \otimes A^{op}) \diamond H$ as bialgebroid:

Theorem 3.2 *Let R be as in Proposition 3.1 and assume that R is moreover a bialgebroid over A , u is a morphism of bialgebroids (with identity map at the level of base algebras) and v is a morphism of bialgebroids (with canonical inclusion $k \rightarrow A$ at the level of base algebras). Then ω is a morphism of bialgebroids (with identity map at the level of base algebras).*

The morphism of bialgebroids $(A \otimes A^{op}) \diamond H \rightarrow A \odot H \odot A$ from Theorem 2.5 can be reobtained easily using this universal property.

References

- [1] P. Bieliavsky, P. Bonneau, Y. Maeda, Universal deformation formulae, symplectic Lie groups and symmetric spaces, *math.QA/0308189*.
- [2] P. Bieliavsky, P. Bonneau, Y. Maeda, Universal deformation formulae for three-dimensional solvable Lie groups, *math.QA/0308188*.
- [3] G. Böhm, K. Szlachányi, Hopf algebroids with bijective antipodes: axioms, integrals and duals, *J. Algebra* **274** (2004), 708–750.
- [4] P. Bonneau, M. Gerstenhaber, A. Giaquinto, D. Sternheimer, Quantum groups and deformation quantization: Explicit approaches and implicit aspects, *J. Math. Phys.* **45** (2004), 3703–3741.
- [5] P. Bonneau, D. Sternheimer, Topological Hopf algebras, quantum groups and deformation quantization, in "Hopf algebras in noncommutative geometry and physics", 55–70, Lecture Notes in Pure and Appl. Math. **239**, Marcel Dekker, New York, 2005.
- [6] D. Bulacu, F. Panaite, F. Van Oystaeyen, Generalized diagonal crossed products and smash products for quasi-Hopf algebras. Applications, *arXiv:math.QA/0506570*.
- [7] C. Cibils, M. Rosso, Hopf bimodules are modules, *J. Pure Appl. Algebra* **128** (1998), 225–231.
- [8] A. Connes, H. Moscovici, Rankin-Cohen brackets and the Hopf algebra of transverse geometry, *Mosc. Math. J.* **4** (2004), 111–130.
- [9] F. Hausser, F. Nill, Diagonal crossed products by duals of quasi-quantum groups, *Rev. Math. Phys.* **11** (1999), 553–629.
- [10] L. Kadison, Pseudo-Galois extensions and Hopf algebroids, *arXiv:math.QA/0508411*.
- [11] M. Khalkhali, B. Rangipour, Para-Hopf algebroids and their cyclic cohomology, *Lett. Math. Phys.* **70** (2004), 259–272.
- [12] J. -H. Lu, Hopf algebroids and quantum groupoids, *Internat. J. Math.* **7** (1996), 47–70.

- [13] F. Panaite, Hopf bimodules are modules over a diagonal crossed product algebra, *Comm. Algebra* **30** (2002), 4049–4058.
- [14] F. Panaite, F. Van Oystaeyen, L-R-smash product for (quasi) Hopf algebras, arXiv:math.QA/0504386.